

Review

Katrina Vilorio and Natasha J. Hill*

Embracing the complexity of matricellular proteins: the functional and clinical significance of splice variation

DOI 10.1515/bmc-2016-0004

Received February 4, 2016; accepted March 24, 2016

Abstract: Matricellular proteins influence wide-ranging fundamental cellular processes including cell adhesion, migration, growth and differentiation. They achieve this both through interactions with cell surface receptors and regulation of the matrix environment. Many matricellular proteins are also associated with diverse clinical disorders including cancer and diabetes. Alternative splicing is a precisely regulated process that can produce multiple isoforms with variable functions from a single gene. To date, the expression of alternate transcripts for the matricellular family has been reported for only a handful of genes. Here we analyse the evidence for alternative splicing across the matricellular family including the secreted protein acidic and rich in cysteine (SPARC), thrombospondin, tenascin and CCN families. We find that matricellular proteins have double the average number of splice variants per gene, and discuss the types of domain affected by splicing in matricellular proteins. We also review the clinical significance of alternative splicing for three specific matricellular proteins that have been relatively well characterised: osteopontin (OPN), tenascin-C (TNC) and periostin. Embracing the complexity of matricellular splice variants will be important for understanding the sometimes contradictory function of these powerful regulatory proteins, and for their effective clinical application as biomarkers and therapeutic targets.

Keywords: alternative splice variation; matricellular; osteopontin; periostin; tenascin-C.

*Corresponding author: **Natasha J. Hill**, Diabetes and Cardiovascular Research Group, Department of Biomolecular Sciences, Kingston University, Penrhyn Road, Kingston-upon-Thames, KT1 2EE, UK, e-mail: n.hill@kingston.ac.uk

Katrina Vilorio: Diabetes and Cardiovascular Research Group, Department of Biomolecular Sciences, Kingston University, Penrhyn Road, Kingston-upon-Thames, UK

Introduction

The extracellular matrix (ECM) is a protein scaffold that provides structural support to tissues, but is also increasingly recognised for the role it plays in providing context to the cell's physical and chemical environment (1). The matrix activates specific cellular signalling pathways through binding of integrins and discoidin domain receptors (DDR), for example. The synergy between integrin and growth factor signalling pathways further regulates cellular responses (2). In addition, the matrix can sequester cytokines and growth factors to achieve high local concentrations protected from degradation, and regulate the activation and release of growth factors (3). The physical properties of the matrix such as rigidity and tension control mechanosensory pathways essential for correct growth, migration and differentiation (2). The matrix is also capable of remodelling to mediate repair and to regulate cell growth, migration and differentiation (4). The distinct matrix composition of individual tissues is an important variable determining tissue-specific characteristics, but as yet remains relatively undefined.

Matricellular proteins are an important component of the ECM. Examples of matricellular proteins include the secreted protein acidic and rich in cysteine (SPARC) family, thrombospondins, tenascins, osteopontin (OPN), the CCN family and periostin (POSTN). Although distinguished from structural matrix proteins such as collagen they nevertheless play an essential regulatory role, and are typically induced during tissue remodelling and repair as well as in many disease states (5). Matricellular proteins can influence matrix formation. For example, SPARC binds to structural matrix components such as collagen and fibronectin, and SPARC null mice have defective ECM that results in cataracts and allows increased tumour growth (6–8). Collagen fibril formation during the foreign body response is also affected in SPARC null mice, as well as in mice lacking TSP-1 or TSP-2 (9).

Matricellular proteins also modulate cellular responses to extracellular signals (10). Many matricellular proteins interact with both growth factors and integrins to regulate cell growth, adhesion and motility. For example, TSP-1 is a physiologically relevant activator of TGF β : it binds to latent TGF β via specific sequences in the thrombospondin repeat region, and causes a conformational change that activates TGF β (11, 12). TSP-1 can also transactivate EGFR and is involved in the regulation of VEGF, FGF, and PDGF signalling (12). Tenascin-X was also recently identified as an activator of TGF β (13). SPARC similarly regulates signalling by a number of growth factors, both directly and indirectly, including TGF β , VEGF, PDGF, FGF and HGF (14, 15). TGF β in turn regulates the expression of a number of matricellular proteins, including SPARC, CCN1, TSP-1, TSP-2 and POSTN (16–19).

In terms of integrin binding, most matricellular proteins bind to multiple integrin receptors, for example, SPARC, CCN1, OPN, POSTN and fibulin-5 (5). Matricellular proteins can therefore directly regulate cell-matrix interactions with subsequent effects on cell adhesion and migration. The capacity of matricellular proteins to modulate integrin signalling also allows indirect regulation of growth factor signalling through the intersection of these pathways. Although we still have much to learn about the effects of matricellular proteins in regulating cell responses to structural matrix proteins, some interesting illustrations are emerging. For example, TSP-1 binding to fibronectin regulates the interaction of TSP-1 with integrin $\alpha 3\beta 1$ (20). Along similar lines, studies identifying the location of SPARC binding on fibrillar collagen suggest that SPARC may influence the interaction between collagen and integrins, and potentially also collagen interactions with DDR2 proteins (21).

Matricellular proteins such as SPARC and TSP-1 were originally defined by their de-adhesive properties (22), although more recently added members such as OPN and TSP2 are now known to promote cell adhesion (10). Although in many cases the effects of matricellular proteins on cell adhesion relate to integrin binding, the effects of TSP-2 on cell adhesion are thought to be mediated by MMP regulation, though altered matrix and growth factor interactions may also play a role (10). Many matricellular proteins in fact regulate MMPs, for example, the SPOCK family of proteins exhibit a complex regulation of MMP activity (23) and SPARC can upregulate expression of multiple MMPs in cancer cells (24). Through MMP regulation matricellular proteins can further influence not just cell adhesion and migration but also cell growth, as MMPs can activate and release growth factors such as VEGF and TGF β (25). Interestingly, many matricellular proteins are also substrates of MMPs and other proteases, and cleavage

of matricellular proteins can expose new active domains and binding sites, or disrupt existing motifs (26, 27). As an illustration, thrombin cleavage of OPN reveals a cryptic non-RGD epitope for integrin binding (28), and cleavage of SPARC produces fragments with increased affinity for collagen (21). Furthermore, cleavage of Hevin/SPARCL1 by MMP3 generates a 'SPARC-like fragment' associated with neovascularisation in glioma and that may compensate for loss of SPARC expression in some tissues (29, 30). Matricellular proteins also regulate protease activity more widely than just MMPs, for example, CCN1 inhibits ADAMTS4 activity (19). This highly complex web of interactions between different matricellular proteins and other regulatory matrix components reinforces the inter-connectedness underpinning homeostatic regulation of the matrix environment.

Through both interactions with cell surface receptors and regulation of the matrix environment, matricellular proteins are therefore able to influence wide-ranging fundamental cellular processes including cell adhesion, migration, growth and differentiation. There is a correspondingly large body of literature demonstrating the involvement of matricellular proteins in many and varied human disorders such as cancer, diabetes, cardiovascular disease and ocular disorders. For example, OPN is over-expressed in the stroma of a variety of cancers and increases malignancy and metastasis both *in vivo* and *in vitro* (31). High levels of SPARC expression in the tumour-stroma are similarly associated with disease progression in many cancers, including pancreatic cancer (4). However, in other types of cancer, such as ovarian and prostate cancers, SPARC expression is downregulated (32). This complexity is also reflected functionally, since although SPARC is overexpressed by the pancreatic tumour stroma, SPARC has been shown to inhibit pancreatic cancer cell growth *in vitro* (33). Similar contradictions are observed with CCN2: while poor prognosis is associated with low CCN2 expression in breast cancer (34), poor prognosis is associated with increased expression in pancreatic cancer (35) and chondrosarcomas (36).

Increased levels of SPARC are also associated with type 2 diabetes, obesity and insulin resistance, both in serum and in adipose tissue (37). SPARC expression is regulated by metabolic parameters such as insulin and glucose in both adipose tissue and in pancreatic stellate cells (15, 37). Increased expression of SPARC is likely to have negative effects in the pancreas since SPARC inhibits β cell responses to growth factors and inhibits β cell survival (15). However, it has also been suggested that ectopic SPARC expression in β cells can actually promote insulin secretion at high levels of glucose (38), perhaps consistent

with the regulation of insulin release by integrin signaling (39). Therefore while the majority of evidence suggests a pathogenic role for SPARC in diabetes, it remains possible that under some circumstances SPARC may have beneficial effects in β cells. Similarly, TSP-1 expression in islet endothelial cells is important for islet function (40), and metabolic parameters such as high glucose also regulate TSP-1 expression (17). As for SPARC, elevated adipose expression of OPN also occurs in obesity, and is thought to predispose to insulin resistance (41). Interestingly, OPN expression is also increased in type 1 diabetes patients (42). Furthermore, the regulation of TGF- β and other growth factors by SPARC, OPN, TSP-1, and members of the CCN family have all been linked with diabetic complications such as nephropathy and retinopathy (43–46).

The contradictory effects of matricellular proteins can therefore be seen at both the functional and clinical level, as illustrated above for the tissue-specific effects of proteins such as SPARC. The specific molecular mechanisms underlying this context dependence are almost entirely unknown, yet this must be addressed if we are to harness the potential therapeutic value of these powerful regulatory proteins. A number of matricellular proteins are currently involved in clinical trials (5, 47), and it will therefore be important that we better understand the mechanisms behind their controversial effects.

The complexity in both form and function of matricellular proteins is likely to explain many of the controversies and tissue-specific effects. Specific matricellular proteins are frequently observed at a range of different molecular weights, as revealed for example by western blotting. The multiple isoforms of these proteins are likely to result from a number of factors, including proteolytic cleavage as discussed above. Matricellular proteins are also subject to a range of post-translational modifications, including glycosylation, phosphorylation and transglutaminase-mediated crosslinking. Where multiple isoforms with different forms or degrees of post-translational modification are present, these isoforms are likely to have distinct functions. For example, transglutaminase cross-linking of OPN (48) and tissue-specific glycosylation of SPARC both result in differential collagen affinity (21).

Alternative splicing also is also likely to contribute to matricellular protein diversity. However, the expression of alternate transcripts for the matricellular family has been reported for only a handful of genes (5). In this review we will examine for the first time the evidence for alternative splicing across the matricellular family, highlighting specific areas for further study. We also illustrate the clinical significance of alternative splicing for three specific matricellular proteins for which splice variants have been

relatively well-characterised: osteopontin, tenascin-C and periostin.

Overview of splice variation in matricellular proteins

Alternative splicing is a precisely regulated process that can produce multiple isoforms with variable functions from a single gene. Alternative splicing is tissue dependent and is regulated by tissue specific *cis*- and *trans*- splicing activators that influence spliceosome assembly (49). We performed an analysis of the splice variants identified in the ENSEMBL database for 22 matricellular proteins, including the SPARC, thrombospondin, tenascin and CCN families, as well as OPN and POSTN. As shown in Table 1, the number of splice variants listed ranges from one to 26, with a mean of eight variants in the set of matricellular genes analysed ($sd=5.3$, $n=22$). Less than half of these are predicted by ENSEMBL to encode a potentially functional protein that would be not be degraded by cellular quality control mechanisms (100/276 variants for all genes analysed). Many non-coding transcripts have retained introns, and such transcripts are typically restricted to the nucleus and thought to be the result of splicing errors, although functionally relevant cases of transcripts with retained introns have been described (50). However, 17 protein coding variants are identified for *SPOCK3*, and 10 for *SPARCL1/Hevin*. Other members of the SPARC family with relatively high numbers of variants listed are *SPARC* and *SPOCK1*, with five predicted protein-coding variants each. Other matricellular genes of interest in this regard are *tenascin C*, *tenascin XB* and *OPN*, having eight, six and five predicted protein variants, respectively.

Recent estimates of the prevalence of alternatively spliced transcripts based on the ENCODE project suggest that, of the 24 000 genes in the human genome, 92–97% express alternatively spliced variants, with an average of four isoforms per gene and on average 2.5 protein coding variants per gene (51, 52). Matricellular proteins therefore have on average around double the average number of splice variants per gene, and double the number of protein coding variants per gene. As discussed below, there is growing evidence for the importance of splice variants of matricellular genes in human health and disease. However, there is clearly still much work to be done in this area. Of the matricellular genes analysed here, only 72/100 protein coding variants listed in ENSEMBL contain the complete coding sequence (CDS), and this is a particular problem for some members of the SPARC family. For

Table 1: Analysis of splice variants of matricellular genes in the ENSEMBL database.

Matricellular Family	Gene name	Description	Other identifiers	Total number ENSEMBL human variants	Number predicted protein coding variants	Number complete CDS listed	Number complete CDS with high confidence (TSL-1)
SPARC	SPARC	Secreted protein, acidic, cysteine-rich (osteonectin)	ON	9	5	1	1
	SPARCL1	SPARC-like 1 (hevin)	MAST9	10	10	3	1
	SPOCK1	SPARC/osteonectin, cwcv and kazal-like domains proteoglycan (testican) 1	SPOCK, testican-1, TIC1	10	5	2	2
	SPOCK2	SPARC/osteonectin, cwcv and kazal-like domains proteoglycan (testican) 2	KIAA0275, testican-2	9	4	4	2
	SPOCK3	SPARC/osteonectin, cwcv and kazal-like domains proteoglycan (testican) 3	Testican-3	26	17	13	3
	SMOC1	SPARC related modular calcium binding 1	-	5	2	2	2
	SMOC2	SPARC related modular calcium binding 2	SMAP2	6	4	3	2
	FSTL1	Follistatin-like 1	FRP, FSL1	10	4	2	1
Thrombospondin	THBS1	Thrombospondin 1	THBS, THBS-1, TSP, TSP-1, TSP1	8	2	1	1
	THBS2	Thrombospondin 2	TSP2	6	3	2	2
	THBS3	Thrombospondin 3	-	11	4	4	3
	THBS4	Thrombospondin 4	-	7	2	2	1
	COMP	Cartilage oligomeric matrix protein	EDM1, EPD1, MED, PSACH, THBS5	4	3	3	2
Tenascin	TNC	Tenascin C	DFNA56, HXB, MGC167029, TN	13	8	6	3
	TNR	Tenascin R	-	3	3	2	0
	TNN	Tenascin N	TN-W	3	3	3	0
	TNXB	Tenascin XB	TNXB1, TNXB2, TNXBS, XB, XBS	9	6	5	2
Osteopontin	SPP1	Secreted phosphoprotein 1	BNSP, BSPI, ETA-1, OPN	11	5	4	3
CCN	CYR61	Cysteine-rich, angiogenic inducer, 61	CCN1, GIG1, IGFBP10	3	2	2	1
	CTGF	Connective tissue growth factor	CCN2, IGFBP8	1	1	1	1
	NOV	Nephroblastoma overexpressed	CCN3, IGFBP9	2	1	1	1
Periostin	POSTN	Periostin, osteoblast specific factor	OSF-2, periostin, PN	10	6	6	5
	Total			176	100	72	39
	Mean variants/gene			8.0	4.5	3.3	1.8

The number of human variants for the main matricellular proteins listed in the ENSEMBL database (accessed August 2015). The final column indicates the number of variants for which there is a high transcript support level (TSL). TSL1 is the highest support level, defined as when transcripts are supported by at least one non-suspect mRNA. CDS, coding sequence. Other matricellular proteins not listed here include: R-spondins, short fibulins, small leucine rich proteoglycans (SLRPs), autotaxin, pigment epithelium derived factor (PEDF) and plasminogen activator inhibitor-1 (PAI-s).

example, *SPARC* (1/5 complete CDS), *SPARCL1* (3/10 complete CDS) and *SPOCK1* (2/5 complete CDS). Furthermore, less than 40% of the protein-coding variants have a high support level, classified by ENSEMBL as transcript support level 1 (TSL1). TSL1 indicates that the transcript is supported by at least one non-suspect mRNA (53). Transcripts with support lower than TSL1 are typically based only on either one or more EST sequences, or on a suspect mRNA sequence, or are not supported by either an EST or mRNA sequence (TSL5). Further work will therefore be required to identify and validate predicted variants of these matricellular proteins, in particular the *SPARC* family.

One critical question will be to determine the abundance of these matricellular splice variants. There is evidence that many splice variants identified by high throughput methods are at low frequency, and that these low frequency variants are typically not evolutionarily conserved and therefore of questionable functional significance (54). Furthermore, it has been estimated that around 80% of transcripts in human tissue represent the major isoform, suggesting relatively minor roles for alternative splice variants in most cases (50). However, the number of variants identified for matricellular genes suggests that many matricellular genes may be amongst the exceptions. Splicing is also altered in disease states such as cancer, and we discuss the evidence for the clinical importance of splice variants of three matricellular genes below. It will also be important to determine cell specific expression patterns of matricellular splice variant transcripts and their relative abundance in different tissues, and to gain insight into the factors that regulate their expression. While RNA seq data from the Illumina Body Map project, for example, can be accessed through ENSEMBL and provides tissue specific information on intron-spanning sequencing reads, data for alternative transcript expression abundance in different cell/tissue types is currently not available. Splice variant databases are also not currently linked to high throughput proteomics data such as mass spectrometry. For coding variants where detailed structural and domain functional data is available for the primary isoform it is possible to use bioinformatics tools to make predictions about the function of alternatively spliced proteins, which can then be tested experimentally.

While many of the matricellular splice variants collated in ENSEMBL predict a change in protein sequence, there is also striking splice variation in exons encoding 5'- and 3'-UTR regions of these genes. The potential importance of variation in the 5'- and 3'-UTRs should also not be underestimated, as our understanding grows of how the secondary structure of these UTR can affect

transcript stability, for example, by miRNA-mediated degradation (55), and also regulate the translational machinery (56). In support of the importance of UTR regions in matricellular protein expression, TSP-1 in particular has a long 3' UTR that has been analysed in detail, and is known to contain AU-rich elements that regulate mRNA stability (17). Indeed, the regulation of TSP-1 expression by glucose occurs at the level of translation and is mediated by miR-467 binding to the 3'-UTR (57, 58). *SPARC* expression is also regulated by non-coding RNA, and in particular is regulated by miR029a/b in nasopharyngeal cancer (59). The potential importance of UTR regions should therefore not be overlooked when considering variants affecting UTR rather than coding regions.

Osteopontin splice variants – evidence for association with specific cancer types

Osteopontin is a member of the SIBLING family of glycoproteins and is expressed from the *SPP1* gene on human chromosome 4. As shown in Figure 1, a total of six splice variants were identified across the two databases, and in Table 2 we have integrated the IDs for these splice variants in ENSEMBL, Genbank and the common names used in the literature. Three splice variants of OPN have been known for some time, widely referred to as OPN-a, OPN-b and OPN-c in the literature, and have been relatively well characterised, while OPN-4 and OPN-5 were only very recently identified (60). Variant 201 listed in ENSEMBL has not been previously described and appears to affect exon 7 outside the splicing region, although this variant is listed as TSL-2 and remains to be confirmed. Interestingly, the recently described OPN-5 is now the longest transcript and therefore the reference sequence at NCBI, though this transcript is not listed in ENSEMBL. This unusual variant has an alternate transcriptional start site, and is discussed further below.

The OPN protein is relatively unstructured and lacks predicted domains identified by bioinformatics searches (61, 62). However, the regions involved in ligand binding have been characterised, as indicated in Figure 1A. There are three integrin-binding motifs in the central region of the protein, plus a C-terminal region responsible for CD44 binding and containing an EF-hand calcium binding motif (63). However, the N-terminal region adjacent to the signal peptide that is affected by splicing lacks any known ligand binding domains (64), although this region does include




Figure 1: Alternative splicing of the human osteopontin gene (*SPP-1*) and associated protein domains.

(A) The osteopontin (OPN) protein is relatively unstructured and lacks conserved domains. Instead, the regions involved in ligand binding are indicated. Splicing affects a region in the N-terminal region of the protein following the signal peptide (SP; AA 31-72 in the OPN-a sequence), shown in orange. This region does not contain any known ligand binding domains but does include O-linked glycosylation and phosphorylation sites (26). There are also three integrin-binding motifs in the central region of the protein, plus a C-terminal region responsible for CD44 binding and containing an EF-hand calcium binding motif (63). The site of thrombin cleavage (AA 169 in the OPN-a sequence) is shown, and OPN is also cleaved by MMPs at AA 166, 201 and 210. Heparin binding sites have been identified at AA 151-160 and 276-283 (26, 63). Diagram is not to scale. (B) The ENSEMBL database identifies five protein coding alternative transcripts for the *SPP-1* gene, resulting from the alternative splicing of two exons (blue box). (C) The NCBI Gene database also identifies five alternative protein coding transcripts for *SPP-1*, but identifies an additional variant not in ENSEMBL ('OPN-5') that includes a novel exon upstream of the alternatively spliced exons. This exon contains an alternative translation start site. Little is as yet known about the structure or function of this variant, but if the downstream translation start site is used the protein would be expected to be present intracellularly as it lacks the signal peptide for secretion. The common OPN variant names are indicated in (B) and (C), and Table 3 provides an overview of the different IDs. All databases accessed in November 2015. (D) The OPN-5 transcript contains an upstream open reading frame (uORF) that may serve to regulate expression of the downstream protein coding sequence. The presence of the uORF also implies that the OPN-5 transcript can only produce a protein lacking a signal peptide, as discussed in the text. OPN-a is shown for comparison (transcripts not to scale).

O-linked glycosylation and phosphorylation sites, and may also contain sites of interaction with the ECM (26). According to Anborgh et al., the exon labelled 'exon 5' in Figure 1 contains glutamine residues that undergo cross-linking by transglutaminase (31). This exon is absent in both OPN-c and OPN-4 variants, suggesting that these variants may be unable to form polymeric structures (Table 2 and Figure 1). Cross-linking enhances the ability of OPN to bind collagen (48), suggesting that OPN-c and OPN-4 may have reduced collagen binding. Furthermore, the alternatively spliced exons 5 and 6 contain putative phosphorylation sites (exon numbering as indicated in Figure 1), suggesting that the various splice variants identified will contain distinct phosphorylation patterns (65). The binding of OPN to integrins is known to be regulated by phosphorylation(66), and it has been suggested that OPN-a and OPN-b, but not OPN-c, can activate integrin receptors (67). This would imply that phosphorylation of exon 5 determines integrin binding, though this would need experimental confirmation.

There is also further evidence for functional and clinical diversity of these variants. While RNA seq data in ENSEMBL suggests that OPN-a is expressed in all tissues tested (ENSEMBL accessed March 2016), multiple splice variants of OPN are found in cancer cells. For example, in a study by He et al., OPN-c was detected in breast cancer but not normal tissue, and OPN-c over-expression was found to promote breast cancer cell invasion *in vitro*. In contrast, OPN-a promoted cell adhesion (64). A follow up study with increased number of patient samples found a more complex picture, but demonstrated strong association between nuclear OPN-c and mortality in patients with early onset breast cancer, while expression of OPN-a/b in the cytoplasm was associated with poor prognosis (68). In pancreatic cancer the picture is subtly different. While OPN-a was found to be highly expressed, OPN-c was associated with metastatic disease and OPN-b expression showed a trend towards association with poor survival rates (69). Similarly, in gastric cancer and glioma while OPN-a, OPN-b and OPN-c were all expressed, only OPN-b

Table 2: Osteopontin variants encoded by the *SPP-1* gene.

Common name	ENSEMBL variant	AA	TSL	Genbank ID	Exons present in alternatively spliced region
OPN-a	001	314	1	NM_001040058	5, 6
OPN-b	002	300	1	NM_000582	5
OPN-c	003	287	1	NM_001040060	6
OPN-4	010	^a	2	NM_001251829	
OPN-5	–	–	–	NM_001251830	4, 5, 6
–	201	292	2	–	5, 6

The splice variant identifiers used in the literature, in ENSEMBL and in Genbank are cross-referenced and the alternatively spliced exons in each variant given. The number of predicted amino acids (AA) and the TSL from the ENSEMBL database are also indicated. Matricellular proteins undergo extensive post-translational modification and also proteolytic cleavage, so the observed molecular weight of each variant may not correspond to that predicted by the amino acid sequence. ^aNumber of predicted amino acids not given as coding sequence (CDS) is incomplete at the 3' end in ENSEMBL. All databases accessed in November 2015.

and OPN-c were associated with the severity of disease symptoms, and in particular OPN-c was again found to be associated with metastasis (70–72).

In a study by Tilli et al., a mechanistic analysis helped to shed additional light on the effects of individual isoforms. Stable expression of OPN-a, OPN-b and OPN-c variants in prostate cancer cells revealed that both OPN-b and OPN-c promote xenograft growth, as well as increasing proliferation, migration and invasion. Expression of OPN-b and OPN-c also increased expression of MMP-2, MMP-9 and VEGF (73). In a similar study by Lin et al., stable expression of OPN-b in esophageal carcinoma (EAC) cells increased migration and enhanced invasion compared to cells expressing OPN-c. However, OPN-c expression dramatically reduced cell adhesion, and the authors suggest that OPN isoforms have distinct effects on cell behaviour but can work together to promote EAC progression (60). This is consistent with the clinical evidence described above suggesting that both OPN-b and OPN-c are overexpressed in some cancers, for example pancreatic (69, 74), gastric (70), lung (74), and prostate (73), while OPN-c may be most strongly associated with other cancers, such as breast cancer (64, 74, 75) and ovarian cancer (76).

These and other studies suggest the potential value of OPN isoforms as biomarkers for specific cancers. This idea was tested by Hartung and Weber in a study analysing blood serum from patients with a range of cancers using an OPN isoform specific RT-PCR assay. Increased expression of both OPN-b and OPN-c was observed in the serum of pancreatic cancer patients while OPN-c was elevated in the serum of lung as well as in breast cancers (74). OPN-b was also elevated in the serum of lung cancer patients, and when a cutoff of two SD over normal was used, could detect over 40% lung cancers.

Diabetes is a risk factor for pancreatic cancer, and OPN expression is also increased in diabetes patients (77, 78). Recently, Sarosiek et al. (78) examined the expression of specific OPN splice variants in patients with pancreatic lesions and with diabetes or obesity. In patients with pancreatic lesions, OPN-b was found in 48% of sera, OPN-c in 34%, and both variants in 5% of sera; neither variant was detected in normal sera. However, in patients with pancreatic lesions who also had diabetes and/or obesity, OPN-b was completely absent and the frequency of OPN-c detection was significantly increased. Logistic regression modelling showed that the odds of having diabetes and/or obesity in a patient with OPN-c were seven times higher than patients without OPN-c. This study demonstrates an association between diabetes/obesity and OPN-c expression, and suggests the possibility that OPN splice variants may be involved in the link between pancreatic cancer and diabetes/obesity (78).

OPN-5 was recently described by Lin et al. (60) and is now the longest transcript identified. As mentioned above, this variant has a unique additional exon (exon 4) containing an alternative translation start site. Interestingly, since the signal peptide is encoded by exon 2, the OPN-5 protein created using the downstream translation start site therefore lacks the signal peptide. The Phobius software for predicting signal peptides clearly identifies AA 1-16 of the other OPN variants as a signal peptide, but does not identify a signal peptide sequence in OPN-5 (79), results that are confirmed using the SignalP prediction tool. It is therefore likely that OPN-5 encodes an intracellular isoform. Translation of the OPN-5 transcript using ExpASY indicates that the upstream start codon utilised by all other OPN isoforms creates an upstream open reading frame (uORF) in the OPN-5 transcript, due to the presence of an in-frame stop codon, as illustrated in Figure 1D.

Only the downstream translational start site can therefore create a functional protein. Interestingly, uORF are important regulators of translation, particularly in response to nutritional and metabolic stress (80). The OPN-5 uORF may therefore serve to regulate the translational efficiency of the OPN-5 protein under certain metabolic conditions.

Although matricellular proteins typically contain a signal peptide that targets the protein for secretion, and OPN is generally regarded as a secreted protein, OPN has been detected intracellularly. Nuclear localisation of OPN has been described (81) and an intracellular isoform of osteopontin (OPNi) has also been characterised. This intracellular form is a truncated protein that lacks the N-terminus, including the signal peptide, and is instead localised to the cytoplasm. Interestingly, this protein was not thought to be a splice variant, but instead is thought to arise from the use of an alternate downstream translation start site (82), although other possibilities have also been discussed (28, 83). OPNi has distinct functions to secreted OPN, and in particular plays an important role in innate immune cells, for example dendritic cells and NK cells (82, 83), where it is involved in signalling transduction pathways downstream of innate immune receptors (28). It will therefore be interesting to test whether innate immune cells express the OPN-5 variant.

The second splice variant recently described by Lin et al., OPN-4, is the shortest transcript and lacks all the alternatively spliced exons 4, 5 and 6 (60). This study found that all five characterised isoforms of OPN, including OPN-4 and OPN-5, are co-overexpressed in primary oesophageal cancer (60). However, no other studies have yet examined the function of OPN-4 and OPN-5 or their potential role in disease.

While the predicted molecular weight of the proteins encoded by the five identified OPN splice variants ranges from 30.8 kDa (OPN-4) to 37.2 kDa (OPN-5; predicted molecular weights calculated using ExPasy), the primary observed band in western blotting is typically around 70 kDa, and multiple bands ranging from 41 to 75 kDa are frequently observed (31). However, OPN undergoes extensive post-translational modification, with serine/threonine phosphorylation, glycosylation, tyrosine sulfation and cross-linking, and OPN is also a substrate for thrombin and matrix metalloproteinases (MMPs). The relationship between expression of particular splice variants and observed protein isoforms of different molecular weights is therefore complex and challenging to unravel.

In summary, OPN mediates cell adhesion and regulates signalling by interacting with a range of binding partners such as CD44 and integrins. There is evidence that alternative splicing creates isoforms with distinct

post-translational modifications that are likely to affect the capacity to interact with integrins and collagen. Expression of specific isoforms has been shown to be associated with diseases such as cancer and diabetes, though more recently described variants remain as yet relatively uncharacterised. Understanding the role of these isoforms and their expression has great potential in their use as biomarkers and therapeutic targets.

Tenascin-C splice variants as therapeutic targets

Tenascin-C (TNC) is a large matricellular glycoprotein that forms a 1080–1500 kDa hexamer, and is encoded by the *TNC* gene on human chromosome 9. Like many other matricellular proteins TNC expression is normally limited in the adult but is upregulated during tissue remodelling, for example during wound repair, inflammation and cancer. Indeed, increased expression of TNC by tumour-stroma cells is associated with a wide range of cancers (84). TNC modulates cell adhesion, migration, proliferation and survival (85), and there are ongoing clinical trials in which TNC is targeted for the treatment of various disorders including heart disease, inflammatory disease and cancer (85, 86).


There are 20 human TNC splice variants identified in the ENSEMBL and NCBI Gene databases (see Table 3 for further details), and the exon structures of the subset of variants present in ENSEMBL are shown in Figure 2B as an illustration. As shown in Figure 2A, TNC contains a C-terminal fibrinogen-like globe (FBG) domain and a region towards the N-terminus containing EGF-like repeats. TNC is a low affinity ligand for EGF receptor and binding is mediated by the EGF-like repeats (87). The TNC protein also contains a central domain consisting of fibronectin type III-like (FNIII) repeats that binds various integrin receptors and growth factors (84, 88). Alternative splicing exclusively affects this central FNIII domain and determines the number of FNIII-like repeat units.

There is extensive evidence for functional diversity in alternative splice variants of TNC. This topic has been recently reviewed in detail (89) so only a brief overview is given here. TNC variants are typically divided into ‘long’ and ‘short’ variants, where long TNC variants can contain up to 17 FNIII like repeats. The small TNC variant, TNC-S, (220 kDa) binds to fibronectin, while the large variant, TNC-L, (320 kDa) does not (90). In contrast, it is uniquely the large TNC variants that bind with high affinity to annexin II (91, 92). Furthermore, while short variants

Table 3: Tenascin-C splice variants encoded by the *TNC* gene.

ENSEMBL variant	AA	TSL	Genbank ID	Exons present in alternatively spliced region
TNC-001	2201	1	NM_002160	11–16
TNC-201	1838	5	XM_006717098	11, 16, 17, 20
TNC-011	2019	5	XM_006717098	11, 12, 13, 15, 16, 20
TNC-009	1928	1	XM_006717100	11–15
TNC-007	1838	1	XM_006717098	11, 16, 17, 20
TNC-012	1564	5	XM_005251975	11
–			XM_005251974	11, 20
–			XM_011518629	11,16,20
–			XM_011518626	11, 16, 17, 19, 20
–			XM_011518624	11, 12, 15, 16, 17, 19, 20
–			XM_006717101	11, 12
–			XM_011518628	11–13
–			XM_006717097	11, 12, 13, 15, 16, 19, 20
–			XM_011518623	11, 12, 13, 15, 16, 17, 19, 20
–			XM_005251973	11–14
–			XM_011518625	11–16
–			XM_005251972	11–16, 20
–			XM_011518622	11–17, 20
–			XM_006717096	11–17, 19, 20
–			XM_011518630	11–19

See the legend for Table 2 for further details.

**Figure 2:** Alternative splicing of the human tenascin-C gene (*TNC*) and associated protein domains.

(A) At the protein level, each tenascin-C monomer consists of a signal peptide targeting the protein for secretion (SP), an N terminal tenascin assembly (TA) domain responsible for hexamer formation and a run of 14.5 epidermal growth factor-like (EGF-L) repeats. Towards the C-terminus are between 8 and 17 fibronectin type III-Like (FNIII) repeats, and the number of repeats is determined by alternative splicing; plus a C-terminal Fibrinogen-Like Globe (FBG) domain. The FNIII repeats shown in white are expressed in all variants, while the orange boxes indicate alternatively spliced FNIII repeats. Diagram is not to scale. (B) The ENSEMBL database identifies six alternative protein coding transcripts for the *TNC* gene, resulting from the alternative splicing of nine exons towards the 3' end of the coding sequence (blue box). The original publication by Nies et al. describing the human *TNC* cDNA sequence and domain structure was used to identify the nucleotide sequence corresponding to each named FNIII repeat and identify the corresponding exon in the ENSEMBL database (120). Additional splice variants identified in the NCBI Gene database are given in Table 3. All databases accessed in November 2015.

promote focal adhesion formation and cell adhesion, long variants typically inhibit the formation of focal adhesions and promote cell migration (93). These observations have been important for rationalising previous observations that TNC could either promote or inhibit cell adhesion, and are an excellent illustration of how the identification of alternative splice variants can resolve existing functional controversies.

Interestingly, examination of the splice variants in the Gene database allowed us to identify a previously undescribed variant (XM_011518630) that contains a novel alternatively spliced exon containing a stop codon (exon 18 in Table 3 and Figure 2). This variant contains the alternatively spliced exons 11–17, and therefore would be predicted to encode a protein with a long FNIII-repeat domain, but would lack completely the FBG domain. The FBG domain is important for proteoglycan and integrin binding (94, 95). It has also been suggested that the TNC FBG domain can activate TLR4, although it is not known whether this is a direct effect (96). It will therefore be of interest to further investigate the expression and function of this novel TNC variant.

Physiologically, there is strong evidence that the expression of alternative TNC splice variants is developmentally regulated and tissue-specific (89). Furthermore, RNA seq data from the Illumina Body Map project suggests that the long TNC-001 variant is detected in lung, colon and lymph tissues, while the short TNC-012 variant is detected in breast, ovary and prostate tissue (accessed via ENSEMBL, March 2016). There have also been a large number of studies suggesting that in particular the long TNC isoforms are associated with many different types of cancer, though this may depend on tumour type. A study by Borsi et al. demonstrated that expression of long variants that include the alternatively spliced region was associated with high stromal cellularity (97). More recent studies using antibodies specific for the alternatively spliced regions have shown that TNC variants containing these regions are absent in normal tissue but expressed at the tumour invasion front and are associated with metastasis. For example, a TNC antibody specific to exon 14 detected a high molecular weight isoform in basement membranes and stromal fibroblasts of breast cancer as well as around the vasculature. This isoform was strongly expressed in the stroma of highly invasive breast tissues but was largely undetected in benign and normal tissue (98). Similarly, antibodies with high affinity for the spliced region detected strong expression of large TNC variants in high grade astrocytomas, particularly near vascular structures and proliferating cells (99). There is currently widespread interest in using the tumour

specific expression of TNC variants to target anti-cancer therapies, and growing interest in developing variant-specific TNC antagonists (89).

Functional and clinical significance of periostin splice variants

POSTN is encoded by the *POSTN* gene on human chromosome 13. Similar to OPN and TNC, POSTN binds a number of integrins and regulates cell adhesion and migration (100, 101). POSTN is also a marker of tumour progression in many types of human cancer, including pancreatic, ovarian, and breast carcinomas, and is thought to promote a tumour-supportive microenvironment (100, 102, 103).


Nine *POSTN* splice variants are identified in ENSEMBL and Gene databases, as detailed in Table 4 (the naming of isoforms used by Morra et al. is used throughout the text for simplicity) (18, 104). The POSTN protein includes four fasciclin-like 1 (FAS-1) repeats that mediate integrin binding, as shown in Figure 3. Alternatively splicing exclusively affects the C-terminal region that has no known conserved functional domains and appears to be intrinsically disordered. The functional effect of alternative splicing is therefore difficult to predict. Hoersch and Andrade-Navarro (105) investigated diversity and isoforms of POSTN from an evolutionary perspective, and hypothesised that the C' domain, while lacking in known functional domains, may assume a structure of multiple consecutive β strands when binding to sequential β strand elements of fibronectin domains. This region also contains predicted heparin binding domains and an anomalous nuclear localisation signal (105, 106). The POSTN C-terminal region is also thought to regulate cell–matrix interactions through binding of additional ECM proteins such as collagen, fibronectin and TNC (107). Interestingly, TGF β induced protein (TGF β Ip) is a paralog of POSTN that has the same domain structure except for the absence of this C' region (108–110). Similarly, the *Drosophila* protein midline fasciclin (MFAS) is classified as a POSTN homologue and also lacks this C' region (111). Furthermore, the capacity of POSTN to regulate invasiveness and lung metastasis also maps to the C-terminus of the protein (112, 113), supporting the functional significance of alternative splice variants.

POSTN splice variant expression is regulated during development in multiple tissues, with additional isoforms expressed during foetal development in both lung and kidney (104). Tissue-specific expression has also been

Table 4: Periostin splice variants encoded by the *POSTN* gene.

Common name [Morra and Moch (18) 2011]	Common name [Kim et al. (113) 2008]	ENSEMBL variant	AA	TSL	Genbank ID	Exons present in alternatively spliced region
'Isoform 1'	'WT'	POSTN-001	836	1	NM_006475	17–21
'Isoform 3'	'Variant II'	POSTN-201	781	1	NM_001136935	18, 19, 20
		POSTN-003	809	1	NM_001286665	18–21
'Isoform 7'		POSTN-202	749	1	NM_001286666	20, 21
'Isoform 2'	'Variant III'	POSTN-002	779	1	NM_001135934	19, 20, 21
'Isoform 5'		POSTN-004	808	5	XM_005266231	17–20
'Isoform 4'	'Variant I'	–			NM_001135936	19, 20
'Isoform 8'		–			NM_001286667	20
'Isoform 6'		–			XM_005266232	17, 19, 20, 21

See the legend for Table 2 for further detail. The common names used in Ref. (18) and Ref. (113) are also given.

**Figure 3:** Alternative splicing of the human periostin gene (*POSTN*) and associated protein domains.

(A) The periostin protein consists of a signal peptide (SP) that targets the protein for secretion, a cysteine-rich EMI domain, and four fasciclin-like 1 (FAS-1) repeats. The second and fourth FAS1 domains contain integrin binding domains (black diamonds). The alternatively spliced C' region (shown in orange) has no known functional domains and appears to be intrinsically disordered. The functional effect of alternative splicing is therefore difficult to predict. However, there are predicted heparin binding domains in the C' tail (blue circle), and also an anomalous nuclear localisation signal (green oval) (105, 106). Diagram is not to scale. (B) The ENSEMBL database identifies six alternative protein coding transcripts for *POSTN*, resulting from the alternative splicing of five exons towards the 3' end of the coding sequence (blue box). Additional splice variants identified in the NCBI Gene database are given in Table 4. All databases accessed in November 2015.

observed, with isoform 6 detected in renal but not lung tissue (104). Multiple isoforms were also identified in normal adult kidney (isoforms 3, 5 and 8). Interestingly, isoform 8 was detected more frequently in renal cell carcinoma than in matched normal tissue (18). Of the alternatively spliced exons 17–21, this isoform contains exon 20 alone. Since only the C-terminal exons 14 to 23 were examined this isoform remains to be fully characterised, and it remains to be shown whether this isoform produces a functional protein. Multiple *POSTN* isoforms were also identified in thyroid carcinoma, but expression was not different to normal tissue (114). In support of highly tissue

specific expression patterns, the RNA seq data in ENSEMBL suggests that while isoform 1 is expressed in breast, ovary, testes and heart tissue, isoform 3 is expressed in adipose, colon, kidney, lymph and prostate, isoform 5 is expressed in the thyroid, while the novel isoform *POSTN-003* is expressed in adrenal tissues (ENSEMBL accessed March 2016).

Although *POSTN* is frequently overexpressed in different cancers, *POSTN* mRNA is downregulated in bladder cancer. Kim et al. (113) showed that normal bladder tissues expressed four different *POSTN* splice variants, corresponding to isoforms 1, 2, 3 and 4. The normal bladder

therefore expresses a distinct splice variant profile to the normal kidney. Importantly, all bladder cancer tissues examined had lost expression of the full length mRNA (isoform 1), and 48% of bladder cancer tissues had lost expression of alternatively spliced versions of POSTN. Instead, isoform 4 was the most commonly expressed POSTN variant in bladder cancer tissues. Functional analysis showed that this variant lacks the capacity to inhibit cell invasion and metastasis that were observed for both isoforms 1 and 3. This suggests that a shift from full length to alternatively splice variants of POSTN may be involved in the metastasis of bladder cancer (113). Isoform 4 contains only exons 19 and 20 of the alternatively spliced exons, and like isoform 8 that is associated with renal cell carcinoma, this isoform may also lack important functional domains.

However, further studies are needed to more fully characterise POSTN splice variants. Of the nine potential splice variants identified in the databases, only four have been fully sequenced and studied in some detail (18). Analysis of the predicted proteins encoded by all variants is currently lacking, and further work is needed to investigate, for example, whether the predicted protein coding POSTN isoforms indeed produce functional proteins, and how binding to collagen and other matricellular components is affected in these isoforms.

Perspectives and conclusions

There are interesting parallels between the regions affected by alternative splicing in POSTN and OPN. Both are regions lacking in conserved functional domains (64, 107). Furthermore, both spliced domains are thought to contain heparin-binding sites that mediate matrix binding (18, 26, 105, 106). This suggests that alternative splicing in these matricellular proteins produces variants with altered matrix binding properties; although this will require experimental testing to substantiate. Interestingly, the correspondence between alternatively spliced exons and unstructured proteins or regions, rather than in a globular domain, is consistent with current thinking in terms of the tolerance of protein structures to accommodate variation. It has been observed that splicing is coupled with protein disorder, as such proteins are naturally less prone to mis-folding and degradation (51). However, this is not exclusively the case, and around 28% alternatively spliced variants have split protein domains (51). For TNC, although the complete structure of this large protein is not known, the structure must allow for variable FNIII repeat numbers. Protein repeats can form either open or closed

structures, and open structures tend to be more tolerant of variable copy number (115), suggesting that the TNC FNIII repeats may form an open structure.

We have identified that the protein encoded by the recently described OPN-5 isoform lacks a signal peptide and is therefore likely to be located intracellularly. Intracellular OPN is important in innate immune cells, yet was previously thought not to occur as a result of alternative splicing (79). It will therefore be important to test the expression of OPN-5 in innate immune cells. Similarly, the study of splice variants of other matricellular proteins may also identify novel intracellular variants, and therefore provide an alternative explanation of the intracellular staining pattern observed for many matricellular proteins (116). Endocytosis from the extracellular environment has been observed for some matricellular proteins and is currently thought to be responsible for the presence of these particular matricellular proteins within the cell. For example, there is evidence that SPARC is endocytosed from the extracellular environment and subsequently present in both cytoplasm and associated with the nuclear matrix (117). A similar mechanism of internalisation into the cytoplasm via an endosomal pathway has also been described for CTGF (CCN2) (118, 119). However, it remains possible that intracellular splice variants are also expressed for these or other matricellular proteins.

For the matricellular proteins OPN, TNC and POSTN, there is clear evidence for tissue-specific expression of multiple splice variants, and growing interest in their clinical significance as both prognostic markers and therapeutic targets. In particular, identification of TNC splice variants with opposing functions has helped to resolve controversy over the role of this protein in cell adhesion. The identification of splice variants for the majority of matricellular proteins in ENSEMBL and Gene databases suggests that further research into potential matricellular splice variants is warranted. However, dissecting the diversity of splice variants and their functional significance is challenging. Often, multiple matricellular isoforms are detected at the protein level using traditional techniques such as western blotting, but it can be difficult to decipher splice variants from isoforms resulting from post-translational modifications and enzymatic cleavage. Furthermore, splice variants themselves are subject to additional post-translational modifications, and may not even be detected by existing antibodies. Proteomics techniques such as mass spectrometry analysed in the context of tissue-specific alternative transcript databases will be required to determine the degree to which splice variation at the transcript level is reflected at the protein level.

In addition, the effect of variants affecting 5'- and 3'-UTR regions are likely to be reflected in changes in protein abundance rather than protein sequence. However, despite these challenges, we suggest that embracing the complexity of matricellular protein splice variation will be an important step towards more fully understanding their function and for effective clinical application of these powerful regulatory proteins.

List of abbreviations

CCN	CYR61, CTGF, NOV protein family
CDS	coding sequence
DDRs	discoidin domain receptors
EAC	esophageal carcinoma
ECM	Extracellular matrix
FAS-1	fasciclin-like 1
FBG	fibrinogen-like globe
FNIII	fibronectin type III-like
MMP	matrix metalloproteinase
OPN	osteopontin
POSTN	periostin
SPARC	secreted protein acidic and rich in cysteine
TGF- β	transforming growth factor- β
TGF β 1p	TGF β induced protein
TNC	tenascin-C
TSL	transcript support level
TSP-1	thrombospondin-1
UTR	untranslated region

References

- Cox TR, Ertler JT. Remodeling and homeostasis of the extracellular matrix: implications for fibrotic diseases and cancer. *Dis Model Mech* 2011; 4: 165–78.
- Kim SH, Turnbull J, Guimond S. Extracellular matrix and cell signalling: the dynamic cooperation of integrin, proteoglycan and growth factor receptor. *J Endocrinol* 2011; 209: 139–51.
- Schultz GS, Wysocki A. Interactions between extracellular matrix and growth factors in wound healing. *Wound Repair Regen* 2009; 17: 153–62.
- Arnold SA, Brekken RA. SPARC: a matricellular regulator of tumorigenesis. *J Cell Commun Signal* 2009; 3: 255–73.
- Murphy-Ullrich JE, Sage EH. Revisiting the matricellular concept. *Matrix Biol* 2014; 37: 1–14.
- Gilmour DT, Lyon GJ, Carlton MB, Sanes JR, Cunningham JM, Anderson JR, Hogan BL, Evans MJ, Colledge WH. Mice deficient for the secreted glycoprotein SPARC/osteonectin/BM40 develop normally but show severe age-onset cataract formation and disruption of the lens. *EMBO J* 1998; 17: 1860–70.
- Brekken RA, Puolakkainen P, Graves DC, Workman G, Lubkin SR, Sage EH. Enhanced growth of tumors in SPARC null mice is associated with changes in the ECM. *J Clin Invest* 2003; 111: 487–95.
- Yan Q, Clark JJ, Wight TN, Sage EH. Alterations in the lens capsule contribute to cataractogenesis in SPARC-null mice. *J Cell Sci* 2002; 115(Pt 13): 2747–56.
- Morris AH, Kyriakides TR. Matricellular proteins and biomaterials. *Matrix Biol* 2014; 37: 183–91.
- Bornstein P, Sage EH. Matricellular proteins: extracellular modulators of cell function. *Curr Opin Cell Biol* 2002; 14: 608–16.
- Schultz-Cherry S, Chen H, Mosher DF, Misenheimer TM, Krutzsch HC, Roberts DD, Murphy-Ullrich JE. Regulation of transforming growth factor- β activation by discrete sequences of thrombospondin 1. *J Biol Chem* 1995; 270: 7304–10.
- Sweetwyne MT, Murphy-Ullrich JE. Thrombospondin1 in tissue repair and fibrosis: TGF- β -dependent and independent mechanisms. *Matrix Biol* 2012; 31: 178–86.
- Alcaraz LB, Exposito JY, Chuvin N, Pommier RM, Cluzel C, Martel S, Sentis S, Bartholin L, Lethias C, Valcourt U. Tenascin-X promotes epithelial-to-mesenchymal transition by activating latent TGF- β . *J Cell Biol* 2014; 205: 409–28.
- Rivera LB, Bradshaw AD, Brekken RA. The regulatory function of SPARC in vascular biology. *Cell Mol Life Sci* 2011; 68: 3165–73.
- Ryall C, Viloría K, Lhaf F, Walker A, King A, Jones P, Mackintosh D, McNeice R, Kocher H, Flodstrom-Tullberg M, Edling C, Hill N. Novel role for matricellular proteins in the regulation of islet β cell survival. *J Biol Chem* 2014; 289: 30614–24.
- Wrana J, Overall C, Sodek J. Regulation of the expression of a secreted acidic protein rich in cysteine (SPARC) in human fibroblasts by transforming growth factor beta. Comparison of transcriptional and post-transcriptional control with fibronectin and type I collagen. *Eur J Biochem* 1991; 197: 519–28.
- Stenina-Adognravi O. Invoking the power of thrombospondins: regulation of thrombospondins expression. *Matrix Biol* 2014; 37: 69–82.
- Morra L, Moch H. Periostin expression and epithelial-mesenchymal transition in cancer: a review and an update. *Virchows Arch* 2011; 459: 465–75.
- Chijiwa M, Mochizuki S, Kimura T, Abe H, Tanaka Y, Fujii Y, Shimizu H, Enomoto H, Toyama Y, Okada Y. CCN1 (Cyr61) is overexpressed in human osteoarthritic cartilage and inhibits ADAMTS-4 (aggrecanase 1) activity. *Arthritis Rheumatol* 2015; 67: 1557–67.
- Rodrigues RG, Guo N, Zhou L, Sipes JM, Williams SB, Templeton NS, Gralnick HR, Roberts DD. Conformational regulation of the fibronectin binding and $\alpha 3 \beta 1$ integrin-mediated adhesive activities of thrombospondin-1. *J Biol Chem* 2001; 276: 27913–22.
- Bradshaw AD. The role of SPARC in extracellular matrix assembly. *J Cell Commun Signal* 2009; 3: 239–46.
- Bornstein P. Thrombospondins as matricellular modulators of cell function. *J Clin Invest* 2001; 107: 929–34.
- Nakada M, Miyamori H, Yamashita J, Sato H. Testican 2 abrogates inhibition of membrane-type matrix metalloproteinases by other Testican family proteins. *Cancer Res* 2003; 63: 3364–9.
- Bradshaw A. Diverse biological functions of the SPARC family of proteins. *Int J Biochem Cell Biol* 2012; 44: 480–8.
- Mott JD, Werb Z. Regulation of matrix biology by matrix metalloproteinases. *Curr Opin Cell Biol* 2004; 16: 558–64.
- Scatena M, Liaw L, Giachelli CM. Osteopontin: a multifunctional molecule regulating chronic inflammation and vascular disease. *Arterioscler Thromb Vasc Biol* 2007; 27: 2302–9.
- Lund SA, Giachelli CM, Scatena M. The role of osteopontin in inflammatory processes. *Cell Commun Signal* 2009; 3: 311–22.

28. Inoue M, Shinohara ML. Intracellular osteopontin (iOPN) and immunity. *Immunol Res* 2011; 49: 160–72.
29. Weaver M, Workman G, Schultz CR, Lemke C, Rempel SA, Sage EH. Proteolysis of matricellular protein Hevin by matrix metalloproteinase-3 produces a SPARC-like fragment (SLF) associated with neovasculature in a murine glioma model. *J Cell Biochem* 2011; 112: 3093–102.
30. Barker TH, Framson P, Puolakkainen PA, Reed M, Funk SE, Sage EH. Matricellular homologs in the foreign body response: hevin suppresses inflammation, but hevin and SPARC together diminish angiogenesis. *Am J Pathol* 2005; 166: 923–33.
31. Anborgh PH, Mutrie JC, Tuck AB, Chambers AF. Pre- and post-translational regulation of osteopontin in cancer. *J Cell Commun Signal* 2011; 5: 111–22.
32. Nagaraju GP, Dontula R, El-Rayes BF, Lakka SS. Molecular mechanisms underlying the divergent roles of SPARC in human carcinogenesis. *Carcinogenesis* 2014; 35: 967–73.
33. Sato N, Fukushima N, Maehara N, Matsubayashi H, Koopman J, Su GH, Hruban RH, Goggins M. SPARC/osteonection is a frequent target for aberrant methylation in pancreatic adenocarcinoma and mediator of tumor-stromal interactions. *Oncogene* 2003; 22: 5021–30.
34. Jiang WG, Watkins G, Fodstad O, Douglas-Jones A, Mokbel K, Mansel RE. Differential expression of the CCN family members Cyr61, CTGF and Nov in human breast cancer. *Endocr Relat Cancer* 2004; 11: 781–91.
35. Hartel M, di Mola FF, Gardini A, Zimmermann A, Di Sebastiano P, Guweidhi A, Innocenti P, Giese T, Giese N, Büchler MW. Desmoplastic reaction influences pancreatic cancer growth behavior. *World J Surg* 2004; 28: 818–25.
36. Shakanaga T, Ozaki T, Ohara N, Asaumi K, Nishida K, Kawai A, Nakanishi T, Takigawa M, Inoue H. Expression of connective tissue growth factor in cartilaginous tumors. *Cancer* 2000; 89: 1466–73.
37. Kos K, Wong S, Tan B, Gummesson A, Jernas M, Franck N, Kerrigan D, Nystrom FH, Carlsson LM, Randeva HS, Pinkney JH, Wilding JP. Regulation of the fibrosis and angiogenesis promoter SPARC/osteonection in human adipose tissue by weight change, leptin, insulin, and glucose. *Diabetes* 2009; 58: 1780–8.
38. Harries LW, McCulloch LJ, Holley JE, Rawling TJ, Welters HJ, Kos K. A role for SPARC in the moderation of human insulin secretion. *PLoS One* 2013; 8: e68253.
39. Kaido T, Yebra M, Cirulli V, Rhodes C, Diaferia G, Montgomery AM. Impact of defined matrix interactions on insulin production by cultured human beta-cells: effect on insulin content, secretion, and gene transcription. *Diabetes* 2006; 55: 2723–9.
40. Olerud J, Mokhtari D, Johansson M, Christoffersson G, Lawler J, Welsh N, Carlsson PO. Thrombospondin-1: an islet endothelial cell signal of importance for beta-cell function. *Diabetes* 2011; 60: 1946–54.
41. Kahles F, Findeisen HM, Bruemmer D. Osteopontin: a novel regulator at the cross roads of inflammation, obesity and diabetes. *Mole Metab* 2014; 3: 384–93.
42. Fierabracci A, Biro PA, Yiangou Y, Mennuni C, Luzzago A, Ludvigsson J, Cortese R, Bottazzo GF. Osteopontin is an autoantigen of the somatostatin cells in human islets: identification by screening random peptide libraries with sera of patients with insulin-dependent diabetes mellitus. *Vaccine* 1999; 18: 342–54.
43. Kos K, Wilding JP. SPARC: a key player in the pathologies associated with obesity and diabetes. *Nat Rev Endocrinol* 2010; 6: 225–35.
44. Nicholas SB, Liu J, Kim J, Ren Y, Collins AR, Nguyen L, Hsueh WA. Critical role for osteopontin in diabetic nephropathy. *Kidney Int* 2010; 77: 588–600.
45. Kong P, Cavalera M, Frangogiannis NG. The role of thrombospondin (TSP)-1 in obesity and diabetes. *Adipocyte* 2014; 3: 81–4.
46. Twigg S, Cooper M. The time has come to target connective tissue growth factor in diabetic complications. *Diabetologia* 2004; 47: 965–8.
47. Sawyer AJ, Kyriakides TR. Matricellular proteins in drug delivery: Therapeutic targets, active agents, and therapeutic localization. *Adv Drug Deliv Rev* 2016; 97: 56–68.
48. Kaartinen MT, Pirhonen A, Linnala-Kankkunen A, Maenpää PH. Cross-linking of osteopontin by tissue transglutaminase increases its collagen binding properties. *J Biol Chem* 1999; 274: 1729–35.
49. Stein S, Lu ZX, Bahrami-Samani E, Park JW, Xing Y. Discover hidden splicing variations by mapping personal transcriptomes to personal genomes. *Nucleic Acids Res* 2015; 43: 10612–22.
50. González-Porta M, Frankish A, Rung J, Harrow J, Brazma A. Transcriptome analysis of human tissues and cell lines reveals one dominant transcript per gene. *Genome Biol* 2013; 14: R70.
51. Hegyi H, Kalmar L, Horvath T, Tompa P. Verification of alternative splicing variants based on domain integrity, truncation length and intrinsic protein disorder. *Nucleic Acids Res* 2011; 39: 1208–19.
52. Tress ML, Martelli PL, Frankish A, Reeves GA, Wesselink JJ, Yeats C, Olason PI, Albrecht M, Hegyi H, Giorgetti A, Raimondo D, Lagarde J, Laskowski RA, Lopez G, Sadowski MI, Watson JD, Fariselli P, Rossi I, Nagy A, Kai W, Stirling Z, Orsini M, Assenov Y, Blankenburg H, Huthmacher C, Ramirez F, Schlicker A, Denoeud F, Jones P, Kerrien S, Orchard S, Antonarakis SE, Reymond A, Birney E, Brunak S, Casadio R, Guigo R, Harrow J, Hermjakob H, Jones DT, Lengauer T, Orengo CA, Patthy L, Thornton JM, Tramontano A, Valencia A. The implications of alternative splicing in the ENCODE protein complement. *Proc Natl Acad Sci USA* 2007; 104: 5495–500.
53. Zerbino DR, Wilder SP, Johnson N, Juettemann T, Flicek PR. The ensembl regulatory build. *Genome Biol* 2015; 16: 56-015-0621-5.
54. Pickrell JK, Pai AA, Gilad Y, Pritchard JK. Noisy splicing drives mRNA isoform diversity in human cells. *PLoS Genet* 2010; 6: e1001236.
55. Valencia-Sanchez MA, Liu J, Hannon GJ, Parker R. Control of translation and mRNA degradation by miRNAs and siRNAs. *Genes Dev* 2006; 20: 515–24.
56. Hughes TA. Regulation of gene expression by alternative untranslated regions. *Trends Genet* 2006; 22: 119–22.
57. Bhattacharyya S, Marinic TE, Krukovets I, Hoppe G, Stenina OI. Cell type-specific post-transcriptional regulation of production of the potent antiangiogenic and proatherogenic protein thrombospondin-1 by high glucose. *J Biol Chem* 2008; 283: 5699–707.
58. Bhattacharyya S, Sul K, Krukovets I, Nestor C, Li J, Adognavi OS. Novel tissue-specific mechanism of regulation of angiogenesis and cancer growth in response to hyperglycemia. *J Am Heart Assoc* 2012; 1: e005967.
59. Qiu F, Sun R, Deng N, Guo T, Cao Y, Yu Y, Wang X, Zou B, Zhang S, Jing T. Mir-29a/b enhances cell migration and invasion in

- nasopharyngeal carcinoma progression by regulating *sparc* and *col3a1* gene expression. *PLoS One* 2015; 10: e0120969.
60. Lin J, Myers AL, Wang Z, Nancarrow DJ, Ferrer-Torres D, Handlogten A, Leverenz K, Bao J, Thomas DG, Wang TD, Orringer MB, Reddy RM, Chang AC, Beer DG, Lin L. Osteopontin (OPN/SPP1) isoforms collectively enhance tumor cell invasion and dissemination in esophageal adenocarcinoma. *Oncotarget* 2015; 6: 22239–57.
 61. Fisher L, Torchia D, Fohr B, Young M, Fedarko N. Flexible structures of SIBLING proteins, bone sialoprotein, and osteopontin. *Biochem Biophys Res Commun* 2001; 280: 460–5.
 62. Mitchell A, Chang HY, Daugherty L, Fraser M, Hunter S, Lopez R, McAnulla C, McMenamin C, Nuka G, Pesseat S, Sangrador-Vegas A, Scheremetjew M, Rato C, Yong SY, Bateman A, Punta M, Attwood TK, Sigrist CJ, Redaschi N, Rivoire C, Xenarios I, Kahn D, Guyot D, Bork P, Letunic I, Gough J, Oates M, Haft D, Huang H, Natale DA, Wu CH, Orengo C, Sillitoe I, Mi H, Thomas PD, Finn RD. The InterPro protein families database: the classification resource after 15 years. *Nucleic Acids Res* 2015; 43(Database issue): D213–21.
 63. O'Regan A, Berman JS. Osteopontin: a key cytokine in cell-mediated and granulomatous inflammation. *Int J Exp Pathol* 2000; 81: 373–90.
 64. He B, Mirza M, Weber G. An osteopontin splice variant induces anchorage independence in human breast cancer cells. *Oncogene* 2006; 25: 2192–202.
 65. Gimba E, Tilli T. Human osteopontin splicing isoforms: known roles, potential clinical applications and activated signaling pathways. *Cancer Lett* 2013; 331: 11–7.
 66. Weber GF, Zawaideh S, Hikita S, Kumar VA, Cantor H, Ashkar S. Phosphorylation-dependent interaction of osteopontin with its receptors regulates macrophage migration and activation. *J Leukoc Biol* 2002; 72: 752–61.
 67. Chae S, Jun H, Lee EG, Yang S, Lee DC, Jung JK, Park KC, Yeom YI, Kim K. Osteopontin splice variants differentially modulate the migratory activity of hepatocellular carcinoma cell lines. *Int J Oncol* 2009; 35: 1409–16.
 68. Zduniak K, Ziolkowski P, Ahlin C, Agrawal A, Agrawal S, Blomqvist C, Fjällskog M, Weber G. Nuclear osteopontin-c is a prognostic breast cancer marker. *Br J Cancer* 2015; 112: 729–38.
 69. Siddiqui AA, Jones E, Andrade D, Shah A, Kowalski TE, Loren DE, Chipitsyna G, Arafat HA. Osteopontin splice variant as a potential marker for metastatic disease in pancreatic adenocarcinoma. *J Gastroenterol Hepatol* 2014; 29: 1321–7.
 70. Tang X, Li J, Yu B, Su L, Yu Y, Yan M, Liu B, Zhu Z. Osteopontin splice variants differentially exert clinicopathological features and biological functions in gastric cancer. *Int J Biol Sci* 2013; 9: 55.
 71. Yan W, Qian C, Zhao P, Zhang J, Shi L, Qian J, Liu N, Fu Z, Kang C, Pu P, You Y. Expression pattern of osteopontin splice variants and its functions on cell apoptosis and invasion in glioma cells. *Neuro Oncol* 2010; 12: 765–75.
 72. Saitoh Y, Kuratsu J, Takeshima H, Yamamoto S, Ushio Y. Expression of osteopontin in human glioma. Its correlation with the malignancy. *Lab Invest* 1995; 72: 55–63.
 73. Tilli TM, Mello KD, Ferreira LB, Matos AR, Accioly MTS, Faria PA, Bellahcène A, Castronovo V, Gimba ER. Both osteopontin-c and osteopontin-b splicing isoforms exert pro-tumorigenic roles in prostate cancer cells. *Prostate* 2012; 72: 1688–99.
 74. Hartung F, Weber GF. RNA blood levels of osteopontin splice variants are cancer markers. *Springerplus* 2013; 2: 110.
 75. Mirza M, Shaughnessy E, Hurley JK, Vanpatten KA, Pestano GA, He B, Weber GF. Osteopontin-c is a selective marker of breast cancer. *Int J Cancer* 2008; 122: 889–97.
 76. Tilli TM, Franco VF, Robbs BK, Wanderley JL, da Silva FR, de Mello KD, Viola JP, Weber GF, Gimba ER. Osteopontin-c splicing isoform contributes to ovarian cancer progression. *Mol Cancer Res* 2011; 9: 280–93.
 77. Huxley R, Ansary-Moghaddam A, De González AB, Barzi F, Woodward M. Type-II diabetes and pancreatic cancer: a meta-analysis of 36 studies. *Br J Cancer* 2005; 92: 2076–83.
 78. Sarosiek K, Jones E, Chipitsyna G, Al-Zoubi M, Kang C, Saxena S, Gandhi AV, Sendiky J, Yeo CJ, Arafat HA. Osteopontin (OPN) isoforms, diabetes, obesity, and cancer; what is one got to do with the other? A new role for OPN. *Journal of Gastrointestinal Surgery* 2015; 19: 639–50.
 79. Phobius [Internet]. Phobius.sbc.su.se. 2015 [cited November 2015]. Available from: <http://Phobius.sbc.su.se>.
 80. Araujo PR, Yoon K, Ko D, Smith AD, Qiao M, Suresh U, Burns SC, Penalva LO. Before it gets started: regulating translation at the 5' UTR. *Comp Funct Genomics* 2012; 12.
 81. Junaid A, Moon MC, Harding GE, Zahradka P. Osteopontin localizes to the nucleus of 293 cells and associates with polo-like kinase-1. *Am J Physiol Cell Physiol* 2007; 292: C919–26.
 82. Shinohara ML, Kim HJ, Kim JH, Garcia VA, Cantor H. Alternative translation of osteopontin generates intracellular and secreted isoforms that mediate distinct biological activities in dendritic cells. *Proc Natl Acad Sci USA* 2008; 105: 7235–39.
 83. Leavenworth JW, Verbinnen B, Wang Q, Shen E, Cantor H. Intracellular osteopontin regulates homeostasis and function of natural killer cells. *Proc Natl Acad Sci USA* 2015; 112: 494–9.
 84. Yoshida T, Akatsuka T, Imanaka-Yoshida K. Tenascin-C and integrins in cancer. *Cell Adh Migr* 2015; 9: 96–104.
 85. Midwood KS, Orend G. The role of tenascin-C in tissue injury and tumorigenesis. *J Cell Commun Signal* 2009; 3: 287–310.
 86. Spenlé C, Saupe F, Midwood K, Burckel H, Noel G, Orend G. Tenascin-C: Exploitation and collateral damage in cancer management. *Cell Adh Migr* 2015; 9: 141–53.
 87. Swindle CS, Tran KT, Johnson TD, Banerjee P, Mayes AM, Griffith L, Wells A. Epidermal growth factor (EGF)-like repeats of human tenascin-C as ligands for EGF receptor. *J Cell Biol* 2001; 154: 459–68.
 88. Chi-Rosso G, Gotwals PJ, Yang J, Ling L, Jiang K, Chao B, Baker DP, Burkly LC, Fawell SE, Koteliensky VE. Fibronectin type III repeats mediate RGD-independent adhesion and signaling through activated β 1 integrins. *J Biol Chem* 1997; 272: 31447–52.
 89. Giblin SP, Midwood KS. Tenascin-C: form versus function. *Cell Adh Migr* 2015; 9: 48–82.
 90. A. Ghert M, Qi W, P. Erickson H, A. Block J, P. Scully S. Tenascin-C splice variant adhesive/anti-adhesive effects on chondrosarcoma cell attachment to fibronectin. *Cell Struct Funct* 2001; 26: 179–87.
 91. Chung CY, Erickson HP. Cell surface annexin II is a high affinity receptor for the alternatively spliced segment of tenascin-C. *J Cell Biol* 1994; 126: 539–48.
 92. Chung CY, Murphy-Ullrich JE, Erickson HP. Mitogenesis, cell migration, and loss of focal adhesions induced by tenascin-C interacting with its cell surface receptor, annexin II. *Mol Biol Cell* 1996; 7: 883–92.
 93. Murphy-Ullrich JE, Lightner VA, Aukhil I, Yan YZ, Erickson HP, Hook M. Focal adhesion integrity is downregulated by the alternatively spliced domain of human tenascin. *J Cell Biol* 1991; 115: 1127–36.

94. Milev P, Fischer D, Häring M, Schulthess T, Margolis RK, Chiquet-Ehrismann R, Margolis RU. The fibrinogen-like globe of tenascin-C mediates its interactions with neurocan and phosphacan/protein-tyrosine phosphatase- ζ/β . *J Biol Chem* 1997; 272: 15501–9.
95. Yokoyama K, Erickson HP, Ikeda Y, Takada Y. Identification of amino acid sequences in fibrinogen gamma-chain and tenascin C C-terminal domains critical for binding to integrin $\alpha V\beta 3$. *J Biol Chem* 2000; 275: 16891–8.
96. Zuliani-Alvarez L, Midwood KS. Fibrinogen-related proteins in tissue repair: how a unique domain with a common structure controls diverse aspects of wound healing. *Adv Wound Care* 2015; 4: 273–85.
97. Borsi L, Carnemolla B, Nicolò G, Spina B, Tanara G, Zardi L. Expression of different tenascin isoforms in normal, hyperplastic and neoplastic human breast tissues. *Int J Cancer* 1992; 52: 688–92.
98. Adams M, Jones JL, Walker RA, Pringle JH, Bell SC. Changes in tenascin-C isoform expression in invasive and preinvasive breast disease. *Cancer Res* 2002; 62: 3289–97.
99. Carnemolla B, Castellani P, Ponassi M, Borsi L, Urbini S, Nicolo G, Dorcaratto A, Viale G, Winter G, Neri D. Identification of a glioblastoma-associated tenascin-C isoform by a high affinity recombinant antibody. *Am J Pathol* 1999; 154: 1345–52.
100. Gillan L, Matei D, Fishman DA, Gerbin CS, Karlan BY, Chang DD. Periostin secreted by epithelial ovarian carcinoma is a ligand for $\alpha V\beta 3$ and $\alpha V\beta 5$ integrins and promotes cell motility. *Cancer Res* 2002; 62: 5358–64.
101. Baril P, Gangeswaran R, Mahon P, Caulee K, Kocher H, Harada T, Zhu M, Kalthoff H, Crnogorac-Jurcovic T, Lemoine N. Periostin promotes invasiveness and resistance of pancreatic cancer cells to hypoxia-induced cell death: role of the $\beta 4$ integrin and the PI3k pathway. *Oncogene* 2007; 26: 2082–94.
102. Erkan M, Kleeff J, Gorbachevski A, Reiser C, Mitkus T, Esposito I, Giese T, Büchler MW, Giese NA, Friess H. Periostin creates a tumor-supportive microenvironment in the pancreas by sustaining fibrogenic stellate cell activity. *Gastroenterology* 2007; 132: 1447–64.
103. Shao R, Bao S, Bai X, Blanchette C, Anderson RM, Dang T, Gishizky ML, Marks JR, Wang XF. Acquired expression of periostin by human breast cancers promotes tumor angiogenesis through up-regulation of vascular endothelial growth factor receptor 2 expression. *Mol Cell Biol* 2004; 24: 3992–4003.
104. Morra L, Rechsteiner M, Casagrande S, von Teichman A, Schraml P, Moch H, Soltermann A. Characterization of periostin isoform pattern in non-small cell lung cancer. *Lung Cancer* 2012; 76: 183–90.
105. Hoersch S, Andrade-Navarro MA. Periostin shows increased evolutionary plasticity in its alternatively spliced region. *BMC Evol Biol* 2010; 10: 30-2148-10-30.
106. Norris RA, Moreno-Rodriguez R, Hoffman S, Markwald RR. The many facets of the matricellular protein periostin during cardiac development, remodeling, and pathophysiology. *J Cell Commun Signal* 2009; 3: 275–86.
107. Norris RA, Damon B, Mironov V, Kasyanov V, Ramamurthi A, Moreno-Rodriguez R, Trusk T, Potts JD, Goodwin RL, Davis J, Hoffman S, Wen X, Sugi Y, Kern CB, Mjaatvedt CH, Turner DK, Oka T, Conway SJ, Molkentin JD, Forgacs G, Markwald RR. Periostin regulates collagen fibrillogenesis and the biomechanical properties of connective tissues. *J Cell Biochem* 2007; 101: 695–711.
108. Skonier J, Neubauer M, Madisen L, Bennett K, Plowman GD, Purchio A. cDNA cloning and sequence analysis of $\beta ig-h3$, a novel gene induced in a human adenocarcinoma cell line after treatment with transforming growth factor- β . *DNA Cell Biol* 1992; 11: 511–22.
109. Ma C, Rong Y, Radloff DR, Datto MB, Centeno B, Bao S, Cheng AW, Lin F, Jiang S, Yeatman TJ, Wang XF. Extracellular matrix protein betaig-h3/TGFBI promotes metastasis of colon cancer by enhancing cell extravasation. *Genes Dev* 2008; 22: 308–21.
110. Horiuchi K, Amizuka N, Takeshita S, Takamatsu H, Katsuura M, Ozawa H, Toyama Y, Bonewald LF, Kudo A. Identification and characterization of a novel protein, periostin, with restricted expression to periosteum and periodontal ligament and increased expression by transforming growth factor β . *Journal of Bone and Mineral Research* 1999; 14: 1239–49.
111. Mosher DF, Johansson MW, Gillis ME, Annis DS. Periostin and TGF- β -induced protein: Two peas in a pod? *Crit Rev Biochem Mol Biol* 2015; 50: 427–39.
112. Kim CJ, Yoshioka N, Tambe Y, Kushima R, Okada Y, Inoue H. Periostin is down-regulated in high grade human bladder cancers and suppresses *in vitro* cell invasiveness and *in vivo* metastasis of cancer cells. *Int J Cancer* 2005; 117: 51–8.
113. Kim CJ, Isono T, Tambe Y, Chano T, Okabe H, Okada Y, Inoue H. Role of alternative splicing of periostin in human bladder carcinogenesis. *Int J Oncol* 2008; 32: 161–9.
114. Bai Y, Nakamura M, Zhou G, Li Y, Liu Z, Ozaki T, Mori I, Kakudo K. Novel isoforms of periostin expressed in the human thyroid. *Jpn Clin Med* 2010; 1: 13.
115. Andrade MA, Petosa C, O'Donoghue SI, Muëller CW, Bork P. Comparison of ARM and HEAT protein repeats. *J Mol Biol* 2001; 309: 1–18.
116. Perbal B. Alternative splicing of CCN mRNAs.... it has been upon us. *J Cell Commun Signal* 2009; 3: 153–7.
117. Gooden M, Vernon R, Bassuk J, Sage E. Cell cycle-dependent nuclear location of the matricellular protein SPARC: association with the nuclear matrix. *J Cell Biochem* 1999; 74: 152–67.
118. Wahab NA, Brinkman H, Mason RM. Uptake and intracellular transport of the connective tissue growth factor: a potential mode of action. *Biochem J* 2001; 359 (Pt 1): 89–97.
119. Kubota S, Hattori T, Shimo T, Nakanishi T, Takigawa M. Novel intracellular effects of human connective tissue growth factor expressed in Cos-7 cells. *FEBS Lett* 2000; 474: 58–62.
120. Nies DE, Hemesath TJ, Kim JH, Gulcher JR, Stefansson K. The complete cDNA sequence of human hexabrachion (tenascin). A multidomain protein containing unique epidermal growth factor repeats. *J Biol Chem* 1991; 266: 2818–23.